




[⁶⁸Ga]Ga-FAPI-04 PET/CT in a patient with endometriosis: a potential game changer?

Caroline Burgard¹ · Florian Rosar¹ · Samer Ezziddin¹ · Bashar H. Hamoud² · Erich-Franz Solomayer² · Matthias W. Laschke³ · Alin Constantin² · Mark Bartholomä¹

Received: 14 February 2025 / Accepted: 27 March 2025
© The Author(s) 2025



A 43-year-old woman with a known history of endometriosis presented with dysmenorrhea, dyschezia, and hematochezia. She had previously undergone a hysterectomy, during which infiltrative endometriosis in the bowel was

identified. To guide further treatment decisions, a magnet resonance imaging (MRI) was performed, confirming bowel involvement and revealing a circumscribed lesion infiltrating the sigmoid colon. Due to persistent symptoms, the patient opted for surgery. For preoperative planning, a fibroblast activation protein inhibitor (FAPI) positron emission tomography/ computed tomography with [⁶⁸Ga]Ga-FAPI-04 was conducted. PET images, acquired 10 min after administration 148 MBq of [⁶⁸Ga]Ga-FAPI-04, showed intense tracer uptake (Fig. A; left column: maximum intensity projection; right column: axial slices of PET/CT fusion and PET; big arrow head: endometriosis in the sigmoid colon, SUVmax 9.8) at the same location as the MRI finding (Fig. B, axial and sagittal slices of contrast enhanced T1-weighted fat saturated sequence), thus confirming the MRI result. Additionally, the PET scan revealed intense uptake in the left

Caroline Burgard, Florian Rosar, Alin Constantin and Mark Bartholomä contributed equally to this work.

✉ Caroline Burgard
caroline.burgard@uks.eu

¹ Department. of Nuclear Medicine, Saarland University – Medical Center, Kirrberger Str. 100, Geb. 50, 66421 Homburg, Germany

² Department of Gynecology and Obstetrics, Saarland University, Homburg, Germany

³ Institute for Clinical and Experimental Surgery, Saarland University, Homburg, Germany

ovary and the ligamentum teres uteri bilaterally (Fig. A, small arrowheads; SUVmax: 11.3, 7.3 and 5.4, respectively) suggesting the presence of additional endometriotic lesions undetected by MRI. The patient underwent a second surgical procedure, which histopathologically confirmed a deep infiltrating endometriosis extragenitalis.

Endometriosis, characterized by the ectopic growth of endometrial-like tissue, is a common gynecological disease affecting approximately 10% of reproductive-age women [1]. It can cause significant symptoms, including dysmenorrhea, dyspareunia, infertility, and gastrointestinal disturbances [2]. In addition to its physical effects, endometriosis can also affect mental health [3]. Thus, endometriosis significantly impact the quality of life of patients [4]. Diagnosis is primarily symptom-based, which can be challenging due to their variety and the heterogenic phenotype of endometriosis overlapping with other diseases. A definitive diagnosis is mostly confirmed through diagnostic laparoscopy. While transvaginal ultrasound is the first-line imaging modality, MRI is often used as a second-line tool for assessing disease extent and aiding in surgical planning [5].

This case highlights the potential role of ^{68}Ga -FAPI PET/CT in detecting endometriotic lesions. Not only did the scan confirm the known lesion, but also identified additional sites of involvement undetected by MRI. Many studies have demonstrated the biodistribution of the ^{68}Ga -labeled FAPI tracer, often showing intense uptake in the uterus [6–8]. There has been a previous case report showing only faint uptake in one endometriotic lesion [9]. To the best of our knowledge, this is the first confirmed case of endometriosis demonstrated on ^{68}Ga -FAPI PET/CT with clear and intensive uptake. Therefore, ^{68}Ga -FAPI PET/CT could represent a promising diagnostic modality for endometriosis, alongside many malignant diseases in which it has proven effective [10–13]. Our observation is supported by a recent preclinical study that found FAP expression in the immune microenvironment of endometriosis from various locations [14].

This case may provide a rationale for future studies systematically analyzing FAP expression in endometriosis across larger patient cohorts, investigating the value of ^{68}Ga -FAPI PET/CT. A prospective study was recently initiated and is currently underway investigating this issue (NCT06792318).

Author contributions Conceptualization: CB, FR and MB; data collection: CB, FR, BHH, AC and E-FS; original draft preparation: CB, FR and MB; review and editing: BHH, AC, E-FS, MWL and SE supervision: E-FS, MWL, and SE. All authors have read and agreed to the submitted version of the manuscript.

Funding Open Access funding enabled and organized by Projekt DEAL.

Data availability The data that support the findings of this study are available from the corresponding author upon reasonable request.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Viganò P, Parazzini F, Somigliana E, Vercellini P. Endometriosis: epidemiology and aetiological factors. *Best Pract Res Clin Obstet Gynaecol.* 2004;18:177–200.
2. Zondervan KT, Becker CM, Missmer SA. Endometriosis. *N Engl J Med.* 2020;382:1244–56.
3. Gambadauro P, Carli V, Hadlaczy G. Depressive symptoms among women with endometriosis: a systematic review and meta-analysis. *Am J Obstet Gynecol.* 2019;220:230–41.
4. Gete DG, Doust J, Mortlock S, Montgomery G, Mishra GD. Impact of endometriosis on women's health-related quality of life: A national prospective cohort study. *Maturitas.* 2023;174:1–7.
5. Quesada J, Härmä K, Reid S, Rao T, Lo G, Yang N, et al. Endometriosis: A multimodal imaging review. *Eur J of Radiol.* 2023;158: 110610.
6. Giesel FL, Kratochwil C, Lindner T, Marschalek MM, Loktev A, Lehnert W, et al. ^{68}Ga -FAPI PET/CT: Biodistribution and Preliminary Dosimetry Estimate of 2 DOTA-Containing FAP-Targeting Agents in Patients with Various Cancers. *J Nucl Med.* 2019;60:386–92.
7. Kessler L, Ferdinandus J, Hirmas N, Zarrad F, Nader M, Kersting D, et al. Pitfalls and Common Findings in ^{68}Ga -FAPI PET: A Pictorial Analysis. *J Nucl Med.* 2022;63:890–6.
8. Zhang X, Song W, Qin C, Song Y, Liu F, Hu F, et al. Uterine Uptake of ^{68}Ga -FAPI-04 in Uterine Pathology and Physiology. *Clin Nucl Med.* 2022;47:7–13.
9. Zou Y, Hu H, Tian R, Wang R. Focal ^{68}Ga -FAPI-04 Uptake in Endometriosis at the Rectouterine Pouch. *Clin Nucl Med.* 2025;
10. Kratochwil C, Flechsig P, Lindner T, Abderrahim L, Altmann A, Mier W, et al. ^{68}Ga -FAPI PET/CT: Tracer Uptake in 28 Different Kinds of Cancer. *J Nucl Med.* 2019;60:801–5.
11. Mori Y, Dendl K, Cardinale J, Kratochwil C, Giesel FL, Haberkorn U. FAPI PET: Fibroblast Activation Protein Inhibitor Use in Oncologic and Nononcologic Disease. *Radiology.* 2023;306: e220749.
12. Dendl K, Koerber SA, Finck R, Mokoala KMG, Staudinger F, Schillings L, et al. ^{68}Ga -FAPI-PET/CT in patients with various gynecological malignancies. *Eur J Nucl Med Mol Imaging.* 2021;48:4089–100.
13. Wegen S, van Heek L, Linde P, Claus K, Akuamo-Boateng D, Baues C, et al. Head-to-Head Comparison of [^{68}Ga]Ga-FAPI-46-PET/CT and [^{18}F]F-FDG-PET/CT for Radiotherapy Planning in Head and Neck Cancer. *Mol Imaging Biol.* 2022;24:986–94.
14. Kellers F, Lützen U, Verburg F, Lebenatus A, Tesch K, Yalcin F, et al. FAP-positive activated fibroblasts are detectable in the microenvironment of endometriosis and correlate with stroma composition and infiltrating CD8+ and CD68+ cells. *Hum Reprod Open.* 2025;hoaf003.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.