

# Exercises for Theoretical physics V

SoSe 2023

Sheet 1

11.04.2023

## Exercise 1 *Quantum harmonic oscillator*

Let us consider a particle of mass  $m$  in one dimension. We define the operator position  $\hat{x}$  and impulsion  $\hat{p}$  such that they obey the commutation relation  $[\hat{x}, \hat{p}] = i\hbar\mathbf{1}$ . The particle is trapped in a harmonic potential such that it is described by the Hamiltonian

$$\hat{H} = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega^2\hat{x}^2 \quad (1)$$

where  $\omega \in \mathbb{R}$ .

- a) Let us define the annihilation and creation operators  $\hat{a} = \frac{1}{\sqrt{2}}\left(\frac{\hat{x}}{l} + i\frac{l\hat{p}}{\hbar}\right)$  and  $\hat{a}^\dagger = \frac{1}{\sqrt{2}}\left(\frac{\hat{x}}{l} - i\frac{l\hat{p}}{\hbar}\right)$ , where  $l = \sqrt{\frac{\hbar}{m\omega}}$ . Show that  $[\hat{a}, \hat{a}^\dagger] = \mathbf{1}$  and that  $\hat{H} = \hbar\omega(\hat{a}^\dagger\hat{a} + \frac{1}{2}\mathbf{1})$

(1 point)

- b) For  $\hat{N} = \hat{a}^\dagger\hat{a}$ , show then that  $[\hat{N}, \hat{a}] = -\hat{a}$  and  $[\hat{N}, \hat{a}^\dagger] = \hat{a}^\dagger$ .

(1 point)

- c) Let us consider the eigenvectors  $|n\rangle$  of the operator  $\hat{N}$  and  $\hat{H}$ , such that  $\hat{N}|n\rangle = n|n\rangle$ , where  $n \in \mathbb{N}$ . Show that if the ground state  $|0\rangle$  is not degenerate, then so are the other states  $|n\rangle$ .

(1 point)

- d) Let us consider the state  $|\psi_0\rangle = \sum_n c_n|n\rangle$ , where  $c_n = e^{-\frac{|\alpha|^2}{2}} \frac{\alpha^n}{\sqrt{n!}}$  and  $\alpha \in \mathbb{C}$ . Show that  $\langle\psi_0|\psi_0\rangle = 1$ . Determine the expectation values  $\langle\hat{x}\rangle$ ,  $\langle\hat{p}\rangle$ ,  $\langle\hat{x}^2\rangle$  and  $\langle\hat{p}^2\rangle$ .

(2 points)

- e) Determine the expression of the evolved state  $|\psi(t)\rangle = \hat{U}(t)|\psi_0\rangle$ . What form takes then  $\langle\hat{x}(t)\rangle$  and  $\langle\hat{p}(t)\rangle$ ?

(2 points)

## Exercise 2 *Conservation equation of quantum probability*

Let us consider a system described by the position- and time-dependent wavefunction  $\psi(\mathbf{r}, t)$ . The behaviour of the wavefunction is governed by the Schrödinger equation

$$i\hbar\frac{\partial}{\partial t}\psi(\mathbf{r}, t) = -\frac{\hbar^2}{2m}\nabla^2\psi(\mathbf{r}, t) + V(\mathbf{r})\psi(\mathbf{r}, t). \quad (2)$$

- a) Defining the quantum probability density  $\rho(\mathbf{r}, t) = |\psi(\mathbf{r}, t)|^2$ , determine the equation of motion of  $\rho$ .

(2 points)

- b) Show that it can be expressed in terms of a conservation equation

$$\frac{\partial}{\partial t}\rho(\mathbf{r}, t) + \nabla \cdot \mathbf{j}(\mathbf{r}, t) = 0. \quad (3)$$

Give the explicit form of the current of probability  $\mathbf{j}$ .

**Hint:** One shall use that  $\nabla \cdot (f\nabla g) = (\nabla f) \cdot (\nabla g) + f\nabla^2 g$ .

(2 points)

### Exercise 3 *Relativistic notations*

- a) Let us define the four-momentum vector  $P^\mu = mu^\mu = (E/c, p_x, p_y, p_z)$ . Show that its pseudo-norm is constant and deduce the relation

$$E^2 = p^2 c^2 + m^2 c^4. \quad (4)$$

Assuming that the mass term is dominant ( $m^2 c^4 \gg p^2 c^2$ ), determine the form of the energy  $E$ .

(2 points)

- b) The Maxwell tensor  $F_{\mu\nu}$  is defined from the four-potential  $A^\mu = (\phi, A_x, A_y, A_z)$  (in Gauss units) as  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ . Show that in the Lorenz gauge ( $\partial_\mu A^\mu = 0$ ), the equation of motion for the electric potential  $\phi$  and the vector potential  $\mathbf{A}$  take the compact form

$$\partial_\mu \partial^\mu A^\nu = \frac{4\pi}{c} j^\nu, \quad (5)$$

where the four-current is defined as  $j^\nu = (\rho c, j_x, j_y, j_z)$ .

**Hint:** In Gauss units, the Maxwell equations take the form

$$\nabla \cdot \mathbf{E} = 4\pi\rho, \quad (6a)$$

$$\nabla \cdot \mathbf{B} = 0, \quad (6b)$$

$$\nabla \times \mathbf{E} = -\frac{1}{c} \frac{\partial}{\partial t} \mathbf{B}, \quad (6c)$$

$$\nabla \times \mathbf{B} = \frac{1}{c} \left( 4\pi \mathbf{j} + \frac{\partial}{\partial t} \mathbf{E} \right). \quad (6d)$$

(2 points)