

Exercises for Theoretical physics IV

WiSe 2022/23

Sheet 10

11.01.2023

Exercise 23 *Particle number fluctuations in the grandcanonical ensemble*
In the grandcanonical ensemble, the expected values of a quantity $A(\{n_{\mathbf{p},s}\})$, which depends on the occupation numbers $n_{\mathbf{p},s}$, are calculated according to the relation

$$\langle A \rangle = \frac{1}{\mathcal{Q}(V, \beta, z)} \sum_{N=0}^{+\infty} z^N \sum_{\substack{\{n_{\mathbf{p},s}\} \\ \sum n_{\mathbf{p},s} = N}} A(\{n_{\mathbf{p},s}\}) e^{-\beta \sum_{\mathbf{p}} \epsilon_{\mathbf{p}} n_{\mathbf{p},s}} \quad (1)$$

The grandcanonical partition function is given here by

$$\mathcal{Q}(V, \beta, z) = \prod_{\mathbf{p},s} [1 \mp z e^{-\beta \epsilon_{\mathbf{p}}}]^{\mp 1} \quad (2)$$

with a minus in the case of Bose-Einstein statistics and a plus for Fermi-Dirac statistics. Here the occupation numbers $n_{\mathbf{p}}$ run through the values $n_{\mathbf{p},s} \in \{0, 1, 2, 3, \dots\}$ for bosons and $n_{\mathbf{p},s} \in \{0, 1\}$ for fermions.

a) First, using Eq.(1), show the validity of the relation

$$\langle n_{\mathbf{p}} \rangle = \sum_{s=-S}^S \langle n_{\mathbf{p},s} \rangle = -\frac{1}{\beta} \frac{\partial}{\partial \epsilon_{\mathbf{p}}} \ln \mathcal{Q} \quad (3)$$

and determine from this the mean occupation numbers

$$\langle n_{\mathbf{p}} \rangle = (2S + 1) \frac{z e^{-\beta \epsilon_{\mathbf{p}}}}{1 \mp z e^{-\beta \epsilon_{\mathbf{p}}}} \quad (4)$$

with a minus for Bose-Einstein statistics and a plus for Fermi-Dirac statistics.

(2 points)

b) Now derive the relation

$$-\frac{1}{\beta} \frac{\partial \langle n_{\mathbf{p}} \rangle}{\partial \epsilon_{\mathbf{p}}} = \langle n_{\mathbf{p}}^2 \rangle - \langle n_{\mathbf{p}} \rangle^2 \quad (5)$$

and show that the following holds for the relative particle number fluctuations

$$\frac{\langle n_{\mathbf{p}}^2 \rangle - \langle n_{\mathbf{p}} \rangle^2}{\langle n_{\mathbf{p}} \rangle^2} = \left[\frac{1}{\langle n_{\mathbf{p}} \rangle} \pm \frac{1}{2S + 1} \right], \quad (6)$$

with a plus for Bose-Einstein statistics and a minus for Fermi-Dirac statistics.

(2 points)

Exercise 24 *Ideal Boltzmann gas in the grandcanonical ensemble*

Let us consider an ideal of gas of N non-interacting particles of mass m with energy

$$E = \sum_{\mathbf{p}} \frac{p^2}{2m} n_{\mathbf{p}}. \quad (7)$$

Furthermore, we also assume that the gas is put in contact with a reservoir at the set temperature T and chemical potential μ .

a) The internal energy of an ideal gas in the grand canonical ensemble reads

$$U = \frac{1}{\mathcal{Q}(V, \beta, z)} \sum_{N=0}^{+\infty} z^N \sum_{\substack{\{n_{\mathbf{p}}\} \\ \sum n_{\mathbf{p}}=N}} E(\{n_{\mathbf{p}}\}) e^{-\beta E}. \quad (8)$$

Show that it can be written in terms of the grand canonical partition function as follows

$$U = -\frac{\partial}{\partial \beta} \ln \mathcal{Q} \quad (9)$$

(1 point)

In the case of an ideal gas the grand canonical partition function is represented by the following form

$$\mathcal{Q} = \sum_{N=0}^{+\infty} \frac{z^N}{N!} V^N \left(\frac{mk_B T}{2\pi\hbar^2} \right)^{3N/2}. \quad (10)$$

Use the definition of the thermal wavelength $\lambda^{-1} = \sqrt{\frac{mk_B T}{2\pi\hbar^2}}$, and show that the partition function is given by

$$\mathcal{Q} = e^{zV/\lambda^3} \quad (11)$$

(1 point)

b) Use Eq(11) and the relation $\frac{PV}{k_B T} = \ln \mathcal{Q}$ to prove that the internal energy can be represented by the equation of state

$$U = \frac{3}{2} PV. \quad (12)$$

(1 point)